



Vanessa Bailey (Jet Propulsion Laboratory, California Institute of Technology)



Jet Propulsion Laboratory Exoplanet Science Initiative symposium

March 26, 2018


# DIRECT IMAGING OF EXOPLANETARY SYSTEMS WITH WFIRST CORONAGRAPH INSTRUMENT (CGI)

© 2018 California Institute of Technology. Government sponsorship acknowledged. The research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. The decision to implement the WFIRST mission will not be finalized until NASA's completion of the National Environmental Policy Act (NEPA) process. This document is being made available for information purposes only.

# WFIRST Scientific Objectives




The full distribution of planets around stars



National Academy of Sciences  
Astronomy & Astrophysics  
Decadal Survey (2010)








# WFIRST-AFTA

- WFIRST uses— **AFTA** (Astrophysics Focused Telescope Asset)
- AFTA is a repurposed **2.4 m** telescope from another government agency
- The AFTA telescope is already built, and sitting in a storage facility
- WFIRST-AFTA includes a coronagraph to image exoplanets
- This was not envisaged by the decadal survey
- Enabled by the 2.4 meter mirror

3/26/2018




Harris Corporation / TJT Photography




# Coronagraphy


- A coronagraph ‘blocks’ light from a host star, enabling light from an exoplanet to enter the detector
- The contrast between a host star and the planets is large



# Coronagraph technology development




# CGI schematic diagram





# CGI Science Filters



$\lambda_1=575$  nm, 10% (annular, 3-9  $\lambda/D$ )

$\lambda_3=760$  nm, 18% (bow-tie / IFS, 3-9  $\lambda/D$ )

$\lambda_2=660$  nm, 18% (bow-tie / IFS, 3-9  $\lambda/D$ )

$\lambda_4=825$  nm, 10% (annular, 3-19  $\lambda/D$ )

# CGI Modes Table

CGI Filters	$\lambda_{\text{center}}$ (nm)	BW	Channel	Masks	Working Angle	Can use w/ linear polarizers	Starlight Suppression Region	Tested before launch?
1	575	10%	Imager	HLC	3-9 $\lambda/D$	Y	360°	Y
2	660	18%	IFS	SPC	3-9 $\lambda/D$		130°	
2	660	18%	Imager	SPC	3-9 $\lambda/D$	Y	130°	
3	760	18%	IFS	SPC	3-9 $\lambda/D$		130°	Y
3	760	18%	Imager	SPC	3-9 $\lambda/D$	Y	130°	
4	825	10%	Imager	HLC	3-9 $\lambda/D$	Y	360°	
4	825	10%	IFS	HLC	3-9 $\lambda/D$		360°	
4	825	10%	IFS	SPC disk	6.5 $\lambda/D$ - 0.8"		360°	
4	825	10%	Imager	SPC disk	6.5-20 $\lambda/D$	Y	360°	Y

$\lambda_1=575$  nm, 10% (annular, 3-9  $\lambda/D$ )

$\lambda_3=760$  nm, 18% (bow-tie / IFS, 3-9  $\lambda/D$ )

$\lambda_2=660$  nm, 18% (bow-tie / IFS, 3-9  $\lambda/D$ )

$\lambda_4=825$  nm, 10% (annular, 3-19  $\lambda/D$ )

# Powers of 10

- Current best coronagraphs reach a contrast ratio of  $10^{6-7}$
- WFIRST CGI performance predictions  $\sim 10^9$
- All technological milestones have been hit ahead of schedule and contrasts of  $10^{8-9}$  has been shown in lab
- WFIRST will test two different types of coronagraphs for both spectroscopy (shaped pupil) and photometry (hybrid Lyot)
- What we need for direct imaging of an exo-Earth to show biomarkers is probably  $10^{10}$
- The Astro 2020 Decadal Survey will look at HabEx and LUVOIR, two mission concepts that might be able to do this





# Science cases for CGI

- Spectroscopy of young self-luminous giant planets
- Imaging & Spectroscopy of reflected light giant planets
- Blind search for gas giants (reflected & emitted light)
- Imaging and spectroscopy of disks
  - Debris
  - Protoplanetary
  - Exozodi
- Download at [http://sites.nationalacademies.org/SSB/CurrentProjects/SSB\\_180659](http://sites.nationalacademies.org/SSB/CurrentProjects/SSB_180659) “View submitted whitepapers”


# Spectra of self-luminous planets

## Beta Pic b, HR 8799 e, 51 Eri b




- New models needed!
- Others possible with IFS + SPC donut?
- **Halp** accretion?

# Reflected light imaging of known RV planets



3/26/2018

ESI - 3/26/18




*B. Nemati exposure time calculator*

13



# Reflected light spectroscopy of RV planets



- Eps Eri ~ 15hr
- 47 Uma ~ 510hr
- HD 114614 ~ 1600hr

# Blind search: 3 months ~ 10 new planets

Savransky 2016





Figure 4: Depth of search (summed completeness assuming one planet per star at each location in the semi-major axis-Radius phase space) for the WFIRST CGI assuming optimal utilization of 3 months of integration time (96 total targets). Giant planets ( $>5$  Earth radii) are found most frequently ( $>50\%$ ) while the CGI is barely sensitive to 2 Earth-radii planets ( $0.5\%$ ). Albedos were assumed to be uniformly 0.2 for planets below 1.4 Earth radii and 0.5 for larger planets.

# Disk Science


- Known disks guaranteed
- Intermediate disks likely







# Exozodi “contaminants”

HLC 575 nm imaging. Nearby sunlike star. 10zodi disk & embedded jovian planet located at 1.6 AU. Flux scale is square-root stretch in units of photoelectrons/s. Simulated exposure time is 2.8 h. (Courtesy of M. Rizzo, N. Zimmerman and the “Haystacks” team). Left/Right = image without/WITH PSF subtraction.



# Protoplanetary disks & protoplanets




# WFIRST CGI Science Yield vs Instrument Performance: Exo-planetary systems



Science \ Contrast	$10^{-9}$	$3 \times 10^{-9}$	$10^{-8}$	$10^{-7}$
Cool EGPs optical spectra	Yes (10+)	A few	No	No
Cool EGPs optical Images	Yes	Yes	Possibly	No
Young EGPs optical spectra	Yes	Yes	Yes	Some
Young EGPs optical images	Yes	Yes	Yes	Some
Exo-Zodi Disks optical images	10 zodis	40 zodis	~100 zodis	1000 zodis

# Project Schedule






# Opportunities with WFIRST

- 25% Guest Observer in 5 year prime mission
- ~100% GO in extended mission
- Guest Investigator calls throughout mission
- All prime survey science teams will be competed in ~2021
- 2020 Decadal Survey will consider a Probe class Starshade







# Formulation Science Working Group

Jeff Kruk GSFC Project Scientist, **Chair**

Jeremy Kasdin Princeton U. CGI Adjutant Scientist, Co-Chair

David Spergel Princeton U. WFI Adjutant Scientist, Co-Chair

## **SCIENCE TEAM PIs**

Olivier Doré JPL Weak Lensing, Redshift Survey

Ryan Foley U. Illinois Supernovae

Scott Gaudi Ohio State U. Microlensing

Jason Kalirai Johns Hopkins U. GO, Milky Way Science

Bruce Macintosh Stanford U. Coronagraph

Saul Perlmutter LBNL Supernovae

James Rhoads Arizona State U. GO, Cosmic Dawn

Brant Robertson UC Santa Cruz GO, Extragalactic Science

Alexander Szalay Johns Hopkins U. GI, Archival Science

Margaret Turnbull SETI Institute Coronagraph

Benjamin Williams U. Washington GO, Nearby Galaxies

## **EX-OFFICIO**

Dominic Benford NASA HQ Program Scientist

Ken Carpenter GSFC Science Center

Roc Cutri Caltech/IPAC Science Center

Jason Rhodes JPL WFIRST Project Scientist

Roeland van der Marel STScI Science Center

## **SCIENCE TEAM DEPUTIES**

Dave Bennett GSFC Microlensing

Chris Hirata Ohio State U. Weak Lensing,  
Redshift Survey

Nikole Lewis STScI Coronagraph

Aki Roberge GSFC Coronagraph

Yun Wang Caltech/IPAC Weak Lensing,  
Redshift Survey

David Weinberg Ohio State U. Weak Lensing,  
Redshift Survey

## **INTERNATIONAL OBSERVERS**


Anthony Boccaletti ESA Representative

Jean Dupuis CSA Representative

Thomas Henning ESA Representative

Toru Yamada JAXA Representative

# Coronagraphy is Challenging




# Microlensing

Microlensing will enable the detection of additional objects from the size of Mars to 30,000 times the mass of our Sun

# Exoplanet Surveys



Ke



WFIRST

Number of planet detections (assuming 1 per star)

0.1 1 10 100 1000 10000



- 2600 planet detections.
- 370 with Earth mass and below.
- Hundreds of free-floating planets.

WFIRST complements  
Kepler, TESS, and  
PLATO.

3/26/2018

ESI - 3/26/18

M. Penny (OSU)

# WFIRST Complements Kepler

